

NON EXISTENCE OF LEVI FLAT HYPERSURFACES WITH POSITIVE NORMAL BUNDLE IN COMPACT KÄHLER MANIFOLDS OF DIMENSION ≥ 3

SÉVERINE BIARD AND ANDREI IORDAN

ABSTRACT. Let X be a compact Kähler manifold of dimension ≥ 3 and L a C^∞ Levi flat hypersurface in X . Then the normal bundle to the Levi foliation does not admit a Hermitian metric with positive curvature along the leaves. This represents an answer to a conjecture of Marco Brunella.

1. INTRODUCTION

A classical theorem of Poincaré-Bendixson [17], [18], [1] states that every leaf of a foliation of the real projective plane accumulates on a compact leaf or on a singularity of the foliation. As a holomorphic foliation $\mathcal{F}$ of codimension 1 of $\mathbb{CP}_n$, $n \geq 2$, does not contain any compact leaf and its singular set $Sing \mathcal{F}$ is not empty, a major problem in foliation theory is the following: can $\mathcal{F}$ contain a leaf F such that $\overline{F} \cap Sing \mathcal{F} = \emptyset$? If this is the case, then there exists a nonempty compact set K called exceptional minimal, invariant by $\mathcal{F}$ and minimal for the inclusion such that $K \cap Sing \mathcal{F} = \emptyset$. The problem of the existence of an exceptional minimal in $\mathbb{CP}_n$, $n \geq 2$ is implicit in [6].

In [7] D. Cerveau proved a dichotomy under the hypothesis of the existence of a holomorphic foliation $\mathcal{F}$ of codimension 1 of $\mathbb{CP}_n$ which admits an exceptional minimal $\mathfrak{M}$: $\mathfrak{M}$ is a real analytic Levi flat hypersurface in $\mathbb{CP}_n$ (i. e. $T(\mathfrak{M}) \cap JT(\mathfrak{M})$ is integrable, where J is the complex structure of $\mathbb{CP}_n$), or there exists $p \in \mathfrak{M}$ such that the leaf through p has a hyperbolic holonomy and the range of the holonomy morphism is a linearisable abelian group. This gave rise to the conjecture of the non-existence of smooth Levi flat hypersurface in $\mathbb{CP}_n$, $n \geq 2$.

The conjecture was proved for $n \geq 3$ by A. Lins Neto [15] for real analytic Levi flat hypersurfaces and by Y.-T. Siu [21] for C^{12} smooth Levi flat hypersurfaces. The methods of proofs for the real analytic case are very different from the smooth case.

A real hypersurface of class C^2 in a complex manifold is Levi flat if its Levi form vanishes or equivalently, it admits a foliation by complex hypersurfaces. We say that a (non-necessarily smooth) real hypersurface L in a complex manifold X is Levi flat if $X \setminus L$ have two pseudoconvex connected components. An example of (non-smooth) Levi flat hypersurface in $\mathbb{CP}_2$ is $L = \{[z_0, z_1, z_2] : |z_1| = |z_2|\}$, where $[z_0, z_1, z_2]$ are homogeneous coordinates in $\mathbb{CP}_2$ (see [10]).

In [12] Iordan and Matthey proved the non existence of Lipschitz Levi flat hypersurfaces in $\mathbb{CP}_n$, $n \geq 3$, which are of Sobolev class W^s , $s > 9/2$. A principal

Date: July, 30, 2014.

Key words and phrases. Levi flat hypersurface, weighted $\bar{\partial}$ -equation.

element of the proof is that the Fubini-Study metric induces a metric of positive curvature on any quotient of the tangent space.

In [3], M. Brunella proved that the normal bundle to the Levi foliation of a closed real analytic Levi flat hypersurface in a compact Kähler manifold of dimension $n \geq 3$ does not admit any Hermitian metric with leafwise positive curvature. The real analytic hypothesis may be relaxed to the assumption of $C^{2,\alpha}$, $0 < \alpha < 1$, such that the Levi foliation extends to a holomorphic foliation in a neighborhood of the hypersurface.

The main step in his proof is to show that the existence of a Hermitian metric with leafwise positive curvature on the normal bundle to the Levi foliation of a compact Levi flat hypersurface L in a Hermitian manifold X , implies that $X \setminus L$ is strongly pseudoconvex, i.e. there exists on $X \setminus L$ an exhaustion function which is strongly plurisubharmonic outside a compact set. This was generalized in [4] for invariant compact subsets of a holomorphic foliation of codimension one. Of course, if X is the complex projective space, then every proper pseudoconvex domain in X is Stein [23].

Brunella stated also the following conjecture [3]: Let X be a compact Kähler manifold of dimension $n \geq 3$ and L a C^∞ compact Levi flat hypersurface in X . Then the normal bundle to the Levi foliation does not admit any Hermitian metric with leafwise positive curvature.

The assumption $n \geq 3$ is necessary in this conjecture (see Example 4.2 of [3]).

In [5] Brunella and Perrone proved that every leaf of a holomorphic foliation $\mathcal{F}$ of codimension one of a projective manifold X of dimension at least 3 and such that $\text{Pic}(X) = \mathbb{Z}$ accumulates on the singular set of the foliation. In this case the normal bundle to the foliation is ample.

In [16], T. Ohsawa considered a C^∞ Levi flat compact hypersurface L in a Kähler manifold X such that the normal bundle to the Levi foliation admits a fiber metric whose curvature is semipositive of rank $\geq k$ on the holomorphic tangent space to the leaves and proved that $X \setminus L$ admits an exhaustion plurisubharmonic function of logarithmic growth which is strictly k -convex. Then, if $\dim X \geq 3$, he proved that there are no Levi flat real analytic hypersurfaces such that the normal bundle to the Levi foliation admits a fiber metric whose curvature is semipositive of rank ≥ 2 on the holomorphic tangent space to L . The same is also true if L is of class C^∞ and the normal bundle to the Levi foliation extends to a holomorphic line bundle [16].

In this paper we solve the above mentioned conjecture of Brunella for compact Kähler manifolds of dimension ≥ 3 . The principal ingredient of the proof is a refinement of the proof of Brunella [3] of the strong pseudoconvexity of $X \setminus L$: we show that there exist a neighborhood U of L and a function v on U vanishing on L , such that $-i\partial\bar{\partial}\ln v \geq c\omega$ on $U \setminus L$, where $c > 0$ and ω is the $(1,1)$ -form associated to the Kähler metric. Then we use the L^2 estimates of Hörmander for the $\bar{\partial}$ -equation [11] and methods developed in [10].

2. PRELIMINARIES

Let X be a n -dimensional Kähler manifold, Ω a domain in X and σ a positive function on Ω . For $\alpha \in \mathbb{R}$ denote

$$L^2_{(p,q)}(\Omega; \sigma^\alpha) = \left\{ f \in L^2_{(p,q)loc}(\Omega) : \int_{\Omega} |f|^2 \sigma^{2\alpha} dV < \infty \right\}$$

endowed with the norm

$$N_\alpha(f) = \left(\int_\Omega |f|^2 \sigma^{2\alpha} dV \right)^{1/2}.$$

We consider the unbounded operator

$$\bar{\partial} = \bar{\partial}_\alpha : L^2_{(p,q)}(\Omega; \sigma^\alpha) \rightarrow L^2_{(p,q+1)}(\Omega; \sigma^\alpha).$$

Definition 1. Let $\alpha \geq 0$ and $f \in L^2_{(p,q)}(\Omega; \sigma^{-\alpha})$. We say that f verifies the moment condition of order α if $\int_\Omega f \wedge h = 0$ for every $\bar{\partial}$ -closed form $h \in L^2_{(n-p,n-q)}(\Omega; \sigma^\alpha)$.

For a subset A of $\partial\Omega$, we will denote by δ_A the geodesic distance to A .

Let Ω be a pseudoconvex domain in $\mathbb{CP}_n$ and $\delta_{\partial\Omega}$ the geodesic distance to the boundary for the Fubini-Study metric. By using the L^2 estimates for the $\bar{\partial}$ -operator of Hörmander with the weight $e^{-\varphi}$, $\varphi = -\alpha \log \delta_{\partial\Omega}$ which is strongly plurisubharmonic by a theorem of Takeuchi [23], Henkin and Iordan proved in [10] the existence and regularity of the $\bar{\partial}$ equation for $\bar{\partial}$ -closed forms in $L^2_{(p,q)}(\Omega; \delta_{\partial\Omega}^{-\alpha})$ verifying the moment condition. This gives the regularity of the $\bar{\partial}$ -operator in pseudoconcave domains with Lipschitz boundary [10] and, by using a method of Siu [21], [22], the non existence of smooth Levi flat hypersurfaces in $\mathbb{CP}_n$, $n \geq 3$ follows (see [12]). These techniques will be used in the 4th paragraph.

We will use also the following theorem of regularity of $\bar{\partial}$ equation of Brinkschulte [2]:

Theorem 1. [2] Let Ω be a relatively compact domain with Lipschitz boundary in a Kähler manifold and set $\delta_{\partial\Omega}$ the geodesic distance to the boundary of Ω . Let $f \in L^2_{(p,q)}(\Omega; \delta_{\partial\Omega}^{-k}) \cap C^k_{(p,q)}(\bar{\Omega}) \cap C^\infty_{(p,q)}(\Omega)$, $q \geq 1$, $k \in \mathbb{N}$ and $u \in L^2_{(p,q-1)}(\Omega; \delta_{\partial\Omega}^{-k})$ such that $\bar{\partial}u = f$ and $\bar{\partial}_{-k}^* u = 0$, where $\bar{\partial}_{-k}^*$ is the Hilbert space adjoint of the unbounded operator $\bar{\partial}_{-k} : L^2_{(p,q-1)}(\Omega; \delta_{\partial\Omega}^{-k}) \rightarrow L^2_{(p,q)}(\Omega; \delta_{\partial\Omega}^{-k})$. Then for k big enough $u \in C^{s(k)}_{(p,q-1)}(\bar{\Omega})$ where $s(k) \underset{k \rightarrow \infty}{\sim} \sqrt{k}$.

3. STRONG PSEUDOCONVEXITY OF THE COMPLEMENT OF A LEVI FLAT HYPERSURFACE

As was mentioned in [3] and in [16] by taking a double covering, we can assume that L is orientable and the complement of a Levi flat hypersurface L has two connected components. This will be always supposed in the sequel and for an open neighborhood U of L we will denote by U^+ and U^- the two connected components of $U \setminus L$.

In [3] Brunella proved that the complement of a closed Levi flat hypersurface in a compact Hermitian manifold of class $C^{2,\alpha}$, $0 < \alpha < 1$, having the property that the Levi foliation extends to a holomorphic foliation in a neighborhood of L and the normal bundle to the Levi foliation admits a C^2 Hermitian metric with leafwise positive curvature is strongly pseudoconvex. The following proposition strenghtens this result:

Proposition 1. Let L be a compact C^3 Levi flat hypersurface in a Hermitian manifold X of dimension $n \geq 2$, such that the normal bundle $\mathcal{N}_L^{1,0}$ to the Levi foliation admits a C^2 Hermitian metric with leafwise positive curvature. Then there exist a neighborhood U of L , $c > 0$ and a positive function $v \in C^2(U)$, vanishing on L such

that $-i\partial\bar{\partial}\ln v \geq c\omega$ on $U \setminus L$, where ω is the $(1,1)$ -form associated to the metric. Moreover, there exists a nonvanishing continuous function g in a neighborhood of L such that $v = g\delta^2$, where δ is the geodesic distance to L .

Proof. Let $z_0 \in L$. There exist holomorphic coordinates $z = (z_1, \dots, z_{n-1}, z_n) = (z', z_n)$ in a neighborhood of z_0 such that the local parametric equations for L are of the form

$$z_j = w_j, \quad j = 1, \dots, n-1, \quad z_n = \varphi(w', t)$$

where φ is of class C^3 on a neighborhood of the origin in $\mathbb{C}^{n-1} \times \mathbb{R}$, holomorphic in w' and $\frac{\partial \varphi}{\partial t}(z_0) \in \mathbb{R}^*$. We consider a C^3 extension $\psi = (\psi_1, \dots, \psi_n)$ of φ on a neighborhood of the origin in $\mathbb{C}^{n-1} \times \mathbb{C}$, $\psi(w', t + is) = (w', \varphi(w', t) + is)$. Then ψ is a C^3 diffeomorphism in a neighborhood of z_0 and holomorphic in w' . It follows that

$$L = \{(z', z_n) : \rho(z', z_n) = 0\}.$$

where $\rho = \text{Im}(\psi^{-1})_n$. We denote $f = (\psi^{-1})_n(z', z_n)$. Since $\bar{\partial}_{z'} f = 0$ on L , there exists an extension $\tilde{f}$ of class C^3 in a neighborhood of z_0 such that $\bar{\partial}\tilde{f}$ vanishes to order greater than 2 on L , i.e. $D^l \bar{\partial}\tilde{f} = 0$ for $|l| \leq 2$ on L .

So there exists an open finite covering $(\widetilde{U}_j)_{j \in J}$ by holomorphic charts of L such that $\widetilde{U}_j \setminus L = \widetilde{U}_j^+ \cup \widetilde{U}_j^-$ such that $U_j = L \cap \widetilde{U}_j = \{z \in \widetilde{U}_j : \text{Im} \tilde{f}_j = 0\}$, where $\bar{\partial}\tilde{f}_j$ vanishes to order greater than 2 on L and the Levi foliation is given on U_j by $\{z \in U_j : \tilde{f}_j(z) = c_j\}$, $c_j \in \mathbb{R}$. Thus $d\tilde{f}_j = \partial\tilde{f}_j$ is a non vanishing section of $\mathcal{N}_L^{1,0}$ on U_j and by shrinking $\widetilde{U}_j$, we may consider that $d\tilde{f}_j \neq 0$ on $\widetilde{U}_j$.

We may suppose that $\mathcal{N}_L^{1,0}$ is represented by a cocycle $\{g_{jk}\}$ of class C^2 subordinated to the covering $(U_j)_{j \in J}$ and there exist closed $(1,0)$ -forms α_j of class C^2 on U_j holomorphic along the leaves such that $T^{1,0}(U_j) = \ker \alpha_j$ for every $j \in J$ and $\alpha_j = g_{jk}\alpha_k$ on $U_j \cap U_k$. So $(\alpha_j)_{j \in J}$ defines a global form α on L with values in $\mathcal{N}_L^{1,0}$ such that locally on U_j we have $\alpha(z) = \alpha_j(z) \otimes \alpha_j^*(z)$ where α_j^* is the dual frame of α_j . In particular we have $\alpha_k^* = g_{jk}\alpha_j^*$.

Let h be a C^2 Hermitian metric with positive leafwise curvature $\Theta_h(\mathcal{N}_L^{1,0})$ on $\mathcal{N}_L^{1,0}$. h is defined on each U_j by a C^2 function $h_j = |\alpha_j^*|^2$ such that $h_k = |g_{jk}|^2 h_j$ on $U_j \cap U_k$.

Since $\alpha_j = \eta_j d\tilde{f}_j$ on U_j for every j , where η_j are nowhere vanishing functions of class C^2 on U_j holomorphic along the leaves and

$$\frac{1}{\eta_k} (d\tilde{f}_k)^* = \frac{1}{\eta_j} g_{jk} (d\tilde{f}_j)^*$$

on $U_j \cap U_k$, it follows that

$$|g_{jk}(z)|^2 = \left| \frac{\eta_j(z)}{\eta_k(z)} \right|^2 = \frac{h_k(z)}{h_j(z)}, \quad z \in U_j \cap U_k.$$

So

$$h_j |\eta_j|^2 (\text{Im} \tilde{f}_j)^2 - h_k |\eta_k|^2 (\text{Im} \tilde{f}_k)^2$$

vanishes to order greater than 2 on $U_j \cap U_k$ and $\left(h_j |\eta_j|^2 \left(\operatorname{Im} \tilde{f}_j\right)^2\right)_{j \in J}$ defines a jet of order 2 on L . By Whitney extension theorem there exists a C^2 function v on X such that $v - h_j |\eta_j|^2 \left(\operatorname{Im} \tilde{f}_j\right)^2$ vanishes to order 2 on U_j for every $j \in J$. Let $\tilde{\eta}_j, \tilde{h}_j$ be C^2 extensions of η_j, h_j on $\widetilde{U}_j$ and set $\tilde{\alpha}_j = \tilde{\eta}_j d\tilde{f}_j, \tilde{v} = \tilde{h}_j |\tilde{\eta}_j|^2 \left(\operatorname{Im} \tilde{f}_j\right)^2$.

For $z \in \widetilde{U}_j$ denote $E'_z = \left\{V' \in T_z^{1,0}(X) : \langle \partial \operatorname{Im} \tilde{f}_j, V' \rangle = 0\right\}$ and E''_z the orthogonal of E'_z in $T_z^{1,0}(X)$. Then for every $V \in T_z^{1,0}(X)$ there exists $V' \in E'_z, V'' \in E''_z$ such that $V = V' + V''$. The curvature form $\Theta\left(\mathcal{N}_L^{1,0}\right)$ is represented by $-i\partial\bar{\partial}\ln\left(h_j |\alpha_j|^2\right)$ on U_j , so by shrinking $\widetilde{U}_j$ we may suppose that there exists $\beta > 0$ such that $\left(-i\partial\bar{\partial}\ln\left(\tilde{h}_j |\tilde{\alpha}_j|^2\right)\right)(V', \overline{V'}) \geq \beta\omega(V', \overline{V'})$ for every $z \in \widetilde{U}_j$ and $V \in T_z^{1,0}(X)$.

On $\widetilde{U}_j \setminus L$ we have

$$\begin{aligned} (3.1) \quad -i\partial\bar{\partial}\ln \tilde{v} &= -i\partial\bar{\partial}\ln \left(\tilde{h}_j \left| \frac{\tilde{\alpha}_j}{d\tilde{f}_j} \right|^2 \left(\operatorname{Im} \tilde{f}_j\right)^2 \right) \\ &= -i\partial\bar{\partial}\ln \tilde{h}_j |\tilde{\alpha}_j|^2 + i\partial\bar{\partial}\ln \left| d\tilde{f}_j \right|^2 - i\partial\bar{\partial}\ln \left(\operatorname{Im} \tilde{f}_j\right)^2. \end{aligned}$$

Let $z \in \widetilde{U}_j$ and $V \in T_z^{1,0}(X)$. Then $V = V' + V'', V' \in E'_z$ and $V'' \in E''_z$ and

$$\begin{aligned} -i\partial\bar{\partial}\ln \tilde{h}_j |\tilde{\alpha}_j|^2 (V, \overline{V}) &= \left(-i\partial\bar{\partial}\ln \left(\tilde{h}_j |\tilde{\alpha}_j|^2\right)\right)(V', \overline{V'}) \\ &\quad + 2\operatorname{Re} \left(-i\partial\bar{\partial}\ln \left(\tilde{h}_j |\tilde{\alpha}_j|^2\right)(V', \overline{V''})\right) \\ &\quad + \left(-i\partial\bar{\partial}\ln \left(\tilde{h}_j |\tilde{\alpha}_j|^2\right)\right)(V'', \overline{V''}) \end{aligned}$$

For every $\varepsilon > 0$

$$2 \left| \operatorname{Re} \left(-i\partial\bar{\partial}\ln \left(\tilde{h}_j |\tilde{\alpha}_j|^2\right)(V', \overline{V''})\right) \right| \leq \varepsilon\omega(V', \overline{V'}) + \frac{1}{\varepsilon}\omega(V'', \overline{V''}),$$

so

$$\begin{aligned} (3.2) \quad -i\partial\bar{\partial}\ln \tilde{h}_j |\tilde{\alpha}_j|^2 (V, \overline{V}) &\geq \beta\omega(V', \overline{V'}) - \varepsilon\omega(V', \overline{V'}) - \frac{1}{\varepsilon}\omega(V'', \overline{V''}) \\ &\quad - \left\| -i\partial\bar{\partial}\ln \tilde{h}_j |\tilde{\alpha}_j|^2 \right\|_{\omega} \omega(V'', \overline{V''}) \end{aligned}$$

Since $\bar{\partial}\tilde{f}_j$ vanishes to order greater than 2 on L , for every $\gamma > 0$ there exists a neighborhood of L such that

$$(3.3) \quad \left| i\partial\bar{\partial}\ln \left| d\tilde{f}_j \right|^2 (V, \overline{V}) \right| \leq \gamma\omega(V, \overline{V})$$

and

$$(3.4) \quad \left| i\partial\bar{\partial}\operatorname{Im} \tilde{f}_j (V, \overline{V}) \right| \leq \gamma \left(\operatorname{Im} \tilde{f}_j\right) \omega(V, \overline{V}).$$

Let $z \in \widetilde{U}_j \setminus L$. By (3.4) it follows that

$$\begin{aligned}
 -i\partial\bar{\partial}\ln\left(\operatorname{Im}\tilde{f}_j\right)^2(V,\overline{V}) &= \left(-2\frac{i\partial\bar{\partial}\operatorname{Im}\tilde{f}_j}{\operatorname{Im}\tilde{f}_j} + 2i\frac{\partial\operatorname{Im}\tilde{f}_j \wedge \bar{\partial}\operatorname{Im}\tilde{f}_j}{\left(\operatorname{Im}\tilde{f}_j\right)^2}\right)(V,\overline{V}) \\
 (3.5) \qquad &\geq -2\gamma\omega(V,\overline{V}) + 2i\frac{\partial\operatorname{Im}\tilde{f}_j \wedge \bar{\partial}\operatorname{Im}\tilde{f}_j}{\left(\operatorname{Im}\tilde{f}_j\right)^2}(V'',\overline{V''}) \\
 &\geq -2\gamma\omega(V,\overline{V}) + \frac{2\inf_U \left\|\partial\operatorname{Im}\tilde{f}_j\right\|_\omega^2}{\left(\operatorname{Im}\tilde{f}_j\right)^2}\omega(V'',\overline{V''}).
 \end{aligned}$$

By using (3.2), (3.3) and (3.5), from (3.1) we obtain

$$\begin{aligned}
 (-i\partial\bar{\partial}\ln\tilde{v})(V,\overline{V}) &\geq (\beta - \varepsilon)\omega(V',\overline{V'}) \\
 &\quad + \left(\frac{2}{\left(\operatorname{Im}\tilde{f}_j\right)^2}\inf_U \left\|\partial\operatorname{Im}\tilde{f}_j\right\|_\omega^2 - \frac{1}{\varepsilon} - \left\|-i\partial\bar{\partial}\ln\tilde{h}_j|\tilde{\alpha}_j|^2\right\|_\omega\right)\omega(V'',\overline{V''}) \\
 &\quad - 2\gamma\omega(V,\overline{V}).
 \end{aligned}$$

By choosing $0 < \varepsilon < \beta$ and by shrinking $\widetilde{U}_j$ such that $\frac{2}{\left(\operatorname{Im}\tilde{f}_j\right)^2}$ is big enough and γ small enough, we obtain that there exists $c > 0$ such that $-i\partial\bar{\partial}\ln\tilde{v} \geq c\omega$ on $\widetilde{U}_j$. Finally, since $v - \tilde{v}$ vanishes to order greater than 2 on L , it follows that there exists a neighborhood U' of L such that $-\ln v$ is strongly plurisubharmonic on $U' \setminus L$. We can now take $U = \{z \in U' : v(z) < \mu\}$ for $\mu > 0$ small enough.

L is a C^3 manifold, so the signed distance function δ is a defining function of class C^3 for L . Since v is of class C^2 on U and vanishes to order greater than 2 on L , we have $v = g\delta^2$ with g continuous in a neighborhood of L .

Suppose that there exists $x \in L$ such that $g(x) = 0$. Then $v = o(\delta^2)$ in a neighborhood of x . But there exists j such that $x \in U_j$ and $v = h_j|\eta_j|^2\left(\operatorname{Im}\tilde{f}_j\right)^2 + o(\delta^2)$. Since $\operatorname{Im}\tilde{f}_j = 0$ and $d\operatorname{Im}\tilde{f}_j \neq 0$ on L it follows that $|\nabla^2 v|(x) \neq 0$. This contradiction shows that $g(x) \neq 0$ on L . $\square$

4. WEIGHTED ESTIMATES FOR THE $\bar{\partial}$ -EQUATION

Proposition 2. *Let L be a compact C^∞ Levi flat hypersurface in a compact Kähler manifold (X, ω) of dimension $n \geq 3$. We suppose that there exists a neighborhood U of L in X and a positive C^2 function v on $\overline{U}$ such that $-i\partial\bar{\partial}\ln v \geq c\omega$ on U , $c > 0$, and $v = g\delta_L^2$ on U , where g is continuous on $\overline{U}$ and $g(z) \neq 0$ for every $z \in \overline{U}$. Consider a $a > 0$ small enough such that $W = \{z \in U : v < a\} \subset U$. Let σ be a C^2 function on W such that $\sigma = v$ in a neighborhood of $\{z \in U : v = a\}$ and $\sigma = v^{-1}$ on a neighborhood of $\{z \in U : v = 0\} \setminus L$. Then:*

1) *Let $f \in L^2_{(p,q)}(W, \sigma^k)$, $q \geq 1$, a $\bar{\partial}$ -closed form. Then for k big enough, there exists $u \in L^2_{(p,q-1)}(W, \sigma^k)$ such that $\bar{\partial}u = f$ and $N_k(u) \leq N_k(f)$, where $N_k(u) \leq N_k(f)$ means $N_k(u) \leq C_k N_k(f)$, with $C_k > 0$ independent of f .*

2) Let $f \in L^2_{(p,q)}(W, \sigma^{-k})$, $q \geq 1$, a $\bar{\partial}$ -closed form verifying the moment condition of order k . Then for k big enough there exists $u \in L^2_{(p,q-1)}(W, \sigma^{-k})$ such that $\bar{\partial}u = f$ and $N_{-k}(u) \preccurlyeq N_{-k}(f)$.

Proof. 1) Let $0 < \varepsilon < a < 1$ such that $W_\varepsilon = \{z \in X : \varepsilon < v < a\} \subset W$. Then $\partial W_\varepsilon = \partial W_\varepsilon^1 \cup \partial W_\varepsilon^2$ where $\partial W_\varepsilon^1 = \{z \in U : v(z) = a\} = \{z \in U : -\ln \sigma(z) = -\ln a\}$ is strongly pseudoconcave and $\partial W_\varepsilon^2 = \{z \in U : v(z) = \varepsilon\} = \{z \in U : -\ln \sigma(z) = -\ln \varepsilon\}$ is strongly pseudoconvex.

By the classical results of Hörmander [11] and Kohn [13], [14] and detailed in [8] the basic estimate holds for (p, q) -forms if and only if the Levi form has at least $n - q$ positive eigenvalues or at least $q + 1$ negative eigenvalues. This implies that the $\bar{\partial}$ -Neumann problem on W_ε satisfies $\frac{1}{2}$ -subelliptic estimates and that the harmonic space $\mathcal{H}_{(p,q)}(W_\varepsilon)$ is finite dimensional.

More generally, in [19], Shaw proved that the harmonic space $\mathcal{H}_{(p,q)}(\Omega)$ where $\Omega = \Omega_1 \setminus \overline{\Omega_2}$, where Ω_1, Ω_2 are pseudoconvex domains with smooth boundary in $\mathbb{C}^n$, $0 \leq p \leq n$, $1 \leq q < n - 1$ is finite dimensional and in [20] she proved that in fact $\mathcal{H}_{(p,q)}(\Omega) = \{0\}$. The same is true in complex manifolds under the hypothesis that there exists strongly plurisubharmonic functions in a neighborhood of the boundaries.

As in the proof of Proposition 3.1 of [19], we consider a weight σ which is equal to $\ln v$ in a neighborhood of ∂W_ε^1 and $-\ln v$ in a neighborhood of ∂W_ε^2 and we obtain that for every $f \in L^2_{(p,q)}(W_\varepsilon, \sigma^k)$, $q \geq 1$ and k big enough

$$N_k(f)^2 \leq C_k \left(N_k(\bar{\partial}f)^2 + N_k(\bar{\partial}_k^* f)^2 \right),$$

where $C_k > 0$ depends only on the smallest positive eigenvalue of $-i\partial\bar{\partial}\ln \sigma$ which is $\geq c$ and independent of ε . By taking $\varepsilon \rightarrow 0$ it follows that for k big enough, there exists $u \in L^2_{(p,q-1)}(W, \sigma^k)$ such that $\bar{\partial}u = f$ and $N_k(u) \preccurlyeq N_k(f)$.

2) The proof of this point is analogous of those of [10], Proposition 5.3 and we give it for the convenience of the reader. By the first point it follows that for k big enough the range $\mathcal{R}_{(p,q)}^k(W)$ of the operator $\bar{\partial}_k : L^2_{(p,q-1)}(W, \sigma^k) \rightarrow L^2_{(p,q)}(W, \sigma^k)$ is closed and we can find a bounded operator $T_{(p,q)}^k : \mathcal{R}_{(p,q)}^k(W) \rightarrow L^2_{(p,q-1)}(W, \sigma^k)$, such that $\bar{\partial}T_{(p,q)}^k f = f$ for every $f \in \mathcal{R}_{(p,q)}^k(W)$.

For every $\varphi \in \mathcal{R}_{(n-p,n-q+1)}^k(W)$, define $\Phi_f(\varphi) = \int_W f \wedge T_{(n-p,n-q+1)}^k \varphi$. Φ_f is a continuous linear form on $\mathcal{R}_{(n-p,n-q+1)}^k(W)$. By the Hahn-Banach theorem, we extend Φ_f as a linear form $\widetilde{\Phi}_f$ on $L^2_{(n-p,n-q+1)}(W, \sigma^k)$ such that $\|\widetilde{\Phi}_f\| = \|\Phi_f\|$. Since $\left(L^2_{(n-p,n-q+1)}(W, \sigma^k) \right)' = L^2_{(p,q-1)}(W, \sigma^{-k})$ by the pairing

$$(\alpha, \beta) = \int_W \alpha \wedge \beta, \quad \alpha \in L^2_{(p,q-1)}(W, \sigma^{-k}), \quad \beta \in L^2_{(n-p,n-q+1)}(W, \sigma^k),$$

there exists $u \in L^2_{(p,q-1)}(W, \sigma^{-k})$ such that

$$\widetilde{\Phi}_f(\varphi) = \int_W u \wedge \varphi$$

for every $\varphi \in L^2_{(n-p,n-q+1)}(W, \sigma^k)$.

Since f verifies the moment condition of order k , $\Phi_f(\varphi) = \int_W f \wedge \psi$ for every $\psi \in L^2_{(n-p, n-q)}(W, \sigma^k)$ such that $\bar{\partial}\psi = \varphi$. In particular

$$\Phi_f(\varphi) = \int_W f \wedge \psi = \int_W u \wedge \varphi = \int_W u \wedge \bar{\partial}\psi$$

for every $\psi \in L^2_{(n-p, n-q)}(W, \sigma^k)$ such that $\bar{\partial}\psi = \varphi$. So $(-1)^{p+q+1} \bar{\partial}u = f$ and the proof is completed. $\square$

Lemma 1. *Under the hypothesis of Proposition 2, let $f \in C^\infty_{(p,q)}(W \setminus L) \cap L^2_{(p,q)}(W, \delta_{\partial W}^{-k-\frac{1}{2}})$ be a $\bar{\partial}$ -closed form, $1 \leq q \leq n-1$. Then, for k big enough, f verifies the moment condition of order k .*

Proof. Let $h \in L^2_{(n-p, n-q)}(W; \sigma^k)$ be a $\bar{\partial}$ -closed form. Since $n-q \geq 1$, by Proposition 2, for k big enough there exists $g \in L^2_{(n-p, n-q-1)}(W; \sigma^k)$ such that $\bar{\partial}g = h$ and $N_k(g) \leq N_k(h)$.

For $\varepsilon > 0$ let $W_\varepsilon = \{z \in W \mid \delta_L(z) \geq \varepsilon\}$ and $f_\varepsilon = \chi_\varepsilon f$, where χ_ε is a C^∞ function on W such that $\chi_\varepsilon \equiv 1$ on $W_{2\varepsilon}$, $\chi_\varepsilon \equiv 0$ on $W \setminus W_\varepsilon$, $0 \leq \chi_\varepsilon \leq 1$, $|D\chi_\varepsilon| \leq \frac{C}{\varepsilon}$, $C > 0$.

Since there exists $a, b > 0$ such that $a\delta_L^2 \leq \sigma \leq b\delta_L^2$ on W ,

$$\begin{aligned} N_{-k}(\bar{\partial}f_\varepsilon) &= N_{-k}(\bar{\partial}\chi_\varepsilon \wedge f) = \left(\int_{W_\varepsilon \setminus W_{2\varepsilon}} |\bar{\partial}\chi_\varepsilon \wedge f|^2 \sigma^{-2k} dV \right)^{\frac{1}{2}} \\ &\leq \frac{C}{\varepsilon} \left(\int_{W_\varepsilon \setminus W_{2\varepsilon}} |f|^2 \sigma^{-2k} dV \right)^{\frac{1}{2}} \leq 2C \left(\int_{W_\varepsilon \setminus W_{2\varepsilon}} |f|^2 \sigma^{-2k} \delta_L^{-2} dV \right)^{\frac{1}{2}} \\ &\leq 2\sqrt{b}C \left(\int_{W_\varepsilon \setminus W_{2\varepsilon}} |f|^2 \sigma^{-2k-1} dV \right)^{\frac{1}{2}} \xrightarrow{\varepsilon \rightarrow 0} 0. \end{aligned}$$

We consider an exhaustion $(W_i)_{i \in \mathbb{N}}$ of W such that $g \in L^2_{(n-p, n-q-1)}(\partial W_i; \sigma^k)$ and we have

$$\int_W f_\varepsilon \wedge h = \lim_{i \rightarrow \infty} \int_{W_i} f_\varepsilon \wedge \bar{\partial}g = (-1)^{p+q+1} \lim_{i \rightarrow \infty} \left(\int_{W_i} \bar{\partial}f_\varepsilon \wedge g + \int_{\partial W_i} f_\varepsilon \wedge g \right)$$

Since $f = o(\delta_{\partial W}^k)$ on ∂W , it follows that

$$\int_W f_\varepsilon \wedge h = \int_W \bar{\partial}f_\varepsilon \wedge g.$$

So

$$\begin{aligned} \left| \int_W f \wedge h \right| &= \lim_{\varepsilon \rightarrow 0} \left| \int_W f_\varepsilon \wedge \bar{\partial}g \right| = \lim_{\varepsilon \rightarrow 0} \left| \int_W \bar{\partial}f_\varepsilon \wedge g \right| \\ &= \lim_{\varepsilon \rightarrow 0} \left| \int_W \sigma^{-k} \bar{\partial}f_\varepsilon \wedge \sigma^k g \right| \leq \lim_{\varepsilon \rightarrow 0} N_{-k}(\bar{\partial}f_\varepsilon) N_k(g) = 0. \end{aligned}$$

$\square$

Corollary 1. *Let L be a compact C^∞ Levi flat hypersurface in a Kähler manifold X of dimension $n \geq 3$ such that the normal bundle $\mathcal{N}_L^{1,0}$ to the Levi foliation admits a C^2 Hermitian with leafwise positive curvature. Then for k big enough there exists a neighborhood W of L such that for every $\bar{\partial}$ -closed $f \in C^\infty_{(p,q)}(W \setminus L) \cap$*

$L^2_{(p,q)}(W \setminus L, \delta_{\partial W}^{-k})$ $1 \leq q \leq n-1$, there exists $u \in L^2_{(p,q-1)}(W \setminus L, \delta_{\partial W}^{-k})$ such that $\bar{\partial}u = f$ and $N_{-k}(u) \preccurlyeq N_{-k}(f)$.

Proof. By Proposition 1 there exists a neighborhood U of L , $c > 0$ and a positive function $v \in C^2(\bar{U})$, $v \sim \delta_L^2$, such that $-i\partial\bar{\partial}\ln v \geq c\omega$ on $U \setminus L$. So for every $\alpha \in \mathbb{R}$, $L^2_{(p,q)}(U \setminus L, \delta_L^\alpha) = L^2_{(p,q)}(U \setminus L, v^{\alpha/2})$. By Lemma 1, f verifies the moment condition of order k for each k big enough and the Corollary follows from Proposition 2. $\square$

From Theorem 1 and Corollary 1 we obtain:

Corollary 2. *Under the hypothesis of Corollary 1, for k big enough, there exist $u_k \in C^{s(k)}_{(p,q-1)}(\bar{W}) \cap L^2_{(p,q-1)}(W, \delta_{\partial W}^{-k})$ such that $\bar{\partial}u_k = f$ where $s(k) \underset{k \rightarrow \infty}{\sim} \sqrt{k}$.*

5. NON EXISTENCE OF LEVI FLAT HYPERSURFACES

Proposition 3. *Let L be a compact C^∞ Levi flat hypersurface in a Kähler manifold X of dimension $n \geq 3$ such that the normal bundle $\mathcal{N}_L^{1,0}$ to the Levi foliation admits a C^2 Hermitian with leafwise positive curvature. Let $u \in C^\infty_{(p,q)}(L)$, $1 \leq q \leq n-2$, such that $\bar{\partial}_b u = 0$. Then for every k big enough there exists a neighborhood W of L and a $\bar{\partial}$ -closed extension $U_k \in C^k_{(p,q)}(W)$ of u .*

Proof. By Proposition 1 there exist a neighborhood U of L , $c > 0$ and positive function $v \in C^2(\bar{U})$ vanishing on L such that $v = g\delta_L^2$ and $-i\partial\bar{\partial}\ln v \geq c\omega$ on $U \setminus L$. Let $\tilde{u} \in C^\infty_{(p,q)}(X)$ an extension of u such that $\bar{\partial}\tilde{u}$ vanishes to infinite order on L and $\text{supp } \tilde{u}$ is relatively compact in U . Then $\bar{\partial}\tilde{u} \in L^2_{(p,q+1)}(U; \delta_{\partial U}^{-k})$ for every $k \in \mathbb{N}$. Since $q+1 \leq n-1$, by Proposition 2, for k big enough, there exist a neighborhood W and $h^\pm \in L^2_{(p,q)}(W^\pm, \delta_{\partial W}^{-k})$, where $W^\pm = \{z \in U : v \gtrless 0\}$, such that $\bar{\partial}h^\pm = \bar{\partial}\tilde{u}$ on $W^\pm$. By Corollary 2 we can choose $h^\pm \in C^{s(k)}_{(p,q)}(\bar{W}^\pm)$, $s(k) \underset{k \rightarrow \infty}{\sim} \sqrt{k}$. This means that for k big enough, the form h defined as $h^\pm$ on $\bar{W}^\pm$ is of class C^k on $\bar{W}$ and vanishes on L . So $\tilde{u} - h^\pm$ is a smooth $\bar{\partial}$ -closed form on W which is an extension of u . $\square$

Theorem 2. *Let X be a compact Kähler manifold of dimension $n \geq 3$ and L a C^∞ compact Levi flat hypersurface. Then the normal bundle to the Levi foliation does not admit any Hermitian metric of class C^2 with leafwise positive curvature.*

Proof. Suppose that the normal bundle $\mathcal{N}$ to the Levi foliation admits a Hermitian metric of class C^2 with leafwise positive curvature. Since $\mathcal{N}$ is topologically trivial, its curvature form $\Theta^\mathcal{N}$ for the Kähler metric of X is d -exact. So there exists a 1-form u of class C^∞ on L such that $du = \Theta^\mathcal{N}$; we may suppose that u is real and $u = u^{0,1} + \overline{u^{0,1}}$, where $u^{0,1}$ is the $(0,1)$ component of u . Since $\Theta^\mathcal{N}$ is a $(1,1)$ -form, it follows that $\bar{\partial}_b u^{0,1} = 0$, where $\bar{\partial}_b$ is the tangential Cauchy-Riemann operator. By Proposition 3 there exists a C^k -extension $U^{0,1}$ of $u^{0,1}$ to a neighborhood of L , $k \geq 2$, such that $\bar{\partial}U^{0,1} = 0$.

From now on, the proof follows [3].

Since $\dim X \geq 3$, $U^{0,1}$ admits a $\bar{\partial}$ -closed extension $\widetilde{U^{0,1}}$ to the full X . Indeed, consider a C^k extension $V^{0,1}$ of $U^{0,1}$ to X . Since $X \setminus L$ is strongly pseudoconvex

by a theorem of Grauert and Riemenschneider [9], $H^{n-2}(X \setminus L, K_X) = 0$. By Serre duality it follows the group of cohomology with compact support $H_c^2(X \setminus L, \mathcal{O}) = 0$. So there exists $G^{0,1}$ with compact support in $X \setminus L$ such that $\bar{\partial}V^{0,1} = \bar{\partial}G^{0,1}$. It follows that $\widetilde{U^{0,1}} = V^{0,1} - G^{0,1}$ is a $\bar{\partial}$ -closed C^k extension $U^{0,1}$ to X .

By Hodge symmetry and Dolbeault isomorphism $H^{0,1}(X, \mathbb{C}) \approx \overline{H^{1,0}(X, \mathbb{C})} \approx \overline{H^0(X, \Omega_X^1)}$, where Ω_X^1 is the sheaf of holomorphic 1-forms on X . So there exists $\eta \in H^0(X, \Omega_X^1)$ and $\Phi \in C^k(X)$ such that $\widetilde{U^{0,1}} = \bar{\eta} + \bar{\partial}\Phi$. It follows that $\Theta^{\mathcal{N}} = i\partial_b\bar{\partial}_b \text{Im } \Phi$ on L and this gives a contradiction at the point of L where $\text{Im } \Phi$ reaches its maximum. $\square$

Acknowledgement 1. *We would like to thank M. Adachi and the anonymous referee for very useful remarks.*

REFERENCES

1. I. Bendixson, *Sur les courbes dfinies par une quation differentielle*, Acta Mathematica **24** (1901), 1–88.
2. J. Brinkschulte, *The $\bar{\partial}$ -problem with support conditions on some weakly pseudoconvex domains*, Ark. fr Mat. **42** (2004), 259–282.
3. M. Brunella, *On the dynamics of codimension one holomorphic foliations with ample normal bundle*, Indiana Univ. Math. J. **57** (2008), 3101–3113.
4. ———, *Codimension one foliations on complex tori*, Ann. Fac. Sci. Toulouse Math. **19** (2010), 405–418.
5. M. Brunella and C. Perrone, *Exceptional singularities of codimension one holomorphic foliations*, Publ. Mat. **55** (2011), 295–312.
6. C. Camacho, A. Lins Neto and P. Sad, *Minimal sets of foliations in complex projective space*, Publ. Math. de I.H.E.S. **68** (1988), 187–203.
7. D. Cerveau, *Minimaux des feuilletages algébriques de CP_n* , Ann. Inst. Fourier **43** (1993), 1535–1543.
8. G. B. Folland and J. J. Kohn, *The Neumann problem for the Cauchy-Riemann complex*, no. 75, Princeton Univ. Press, Princeton, N. J., 1972.
9. H. Grauert and O. Riemenschneider, *Kälersche Mannigfaltigkeiten mit hyper-q-konvexen Rand*, vol. A Symposium in Honor of S. Bochner, Princeton 1969, Princeton University Press, Princeton, 1970.
10. G. M. Henkin and A. Iordan, *Regularity of $\bar{\partial}$ on pseudoconcave compacts and applications*, Asian J. Math. **4** (2000), no. 4, 855–884, and Erratum to : Regularity of $\bar{\partial}$ on pseudoconcave compacts and applications by G. M. Henkin and A. Iordan, Asian J. Math., 4, 855–884, 2000.
11. L. Hörmander, *L^2 estimates and existence theorems for the $\bar{\partial}$ operator*, Acta Math. **113** (1965), 89–152.
12. A. Iordan and F. Matthey, *Régularité de l’opérateur $\bar{\partial}$ et théorème de Siu sur la nonexistence d’hypersurfaces Levi-plates dans l’espace projectif complexe CP_n , $n \geq 3$* , C. R. Acad. Sc. Paris **346** (2008), 395–400.
13. J. J. Kohn, *Harmonic integrals on strongly pseudoconvex manifolds I*, Ann. of Math. **78** (1963), no. 2, 112–148.
14. ———, *Global regularity for $\bar{\partial}$ on weakly pseudoconvex manifolds*, T.A.M.S. **181** (1973), 272–292.
15. A. Lins Neto, *A note on projective Levi flats and minimal sets of algebraic foliations*, Ann. Inst. Fourier **49** (1999), 1369–1385.
16. T. Ohsawa, *Nonexistence of certain Levi flat hypersurfaces in Kähler manifolds from the viewpoint of positive normal bundles*, Publ. RIMS Kyoto Univ. **49** (2013), 229–239.
17. H. Poincaré, *Mmoire sur les courbes dfinies par une quation differentielle*, Journal de Math. Pures et Appl. **7** (1881), 375–422.
18. ———, *Mmoire sur les courbes dfinies par une quation differentielle*, Journal de Math. Pures et Appl. **8** (1882), 251–296.
19. M.-C. Shaw, *Global solvability and regularity for $\bar{\partial}$ on an annulus between two weakly pseudoconvex domains*, Trans. Amer. Math. Soc. **291** (1985), no. 1, 255–267.

20. ———, *The closed range property for $\bar{\partial}$ on domains with pseudoconcave boundary*, Complex Analysis, Several Complex Variables and Connections with PDE Theory and Geometry, Springer Basel AG, 2010, pp. 307–320.
21. Y.-T. Siu, *Nonexistence of smooth Levi flat hypersurfaces in complex projective spaces of dimension ≥ 3* , Ann. of Math. **151** (2000), 1217–1243.
22. ———, *$\bar{\partial}$ -regularity for weakly pseudoconvex domains in compact Hermitian symmetric spaces with respect to invariant metrics*, Ann. of Math. **156** (2002), 595–621.
23. A. Takeuchi, *Domaines pseudoconvexes sur les variétés kählériennes*, J. Math. Kyoto Univ. **6** (1967), 323–357.

INSTITUT DE MATHÉMATIQUES, UMR 7586 DU CNRS, CASE 247, UNIVERSITÉ PIERRE ET MARIE-CURIE, 4 PLACE JUSSIEU, 75252 PARIS CEDEX 05, FRANCE
E-mail address: `biard@math.jussieu.fr`

INSTITUT DE MATHÉMATIQUES, UMR 7586 DU CNRS, CASE 247, UNIVERSITÉ PIERRE ET MARIE-CURIE, 4 PLACE JUSSIEU, 75252 PARIS CEDEX 05, FRANCE
E-mail address: `iordan@math.jussieu.fr`